SCIENTIFIC REPORTS

OPEN



Received: 16 September 2016 Accepted: 25 January 2017 Published: 09 March 2017

Diet-induced obesity impairs spermatogenesis: a potential role for autophagy

Yang Mu¹, Wen-jie Yan¹, Tai-lang Yin¹, Yan Zhang², Jie Li¹ & Jing Yang¹

Autophagy is an evolutionarily conserved process that plays a crucial role in maintaining a series of cellular functions. It has been found that autophagy is closely involved in the physiological process of spermatogenesis and the regulation of sperm survival and motility. However, the role of autophagy in high-fat diet (HFD)-induced impaired spermatogenesis remains unknown. This study was designed to investigate the role of autophagy in HFD-induced spermatogenesis deficiency and employed chloroquine (CQ) to inhibit autophagy and rapamycin (RAP) to induce autophagy. 3-methyladenine (3-MA) and CQ were administered via intratesticular injection *in vivo*. The effects of CQ and 3-MA on the parameters of spermatozoa co-cultured with palmitic acid (PA) *in vitro* were also investigated. Human semen samples from obese, subfertile male patients were also collected to examine the level of autophagy. The results suggested that HFD mice subjected to CQ showed improved spermatogenesis. Inhibiting autophagy with CQ improved the decreased fertility of HFD male mice. Moreover, the *in vivo* and *in vitro* results indicated that both CQ and 3-MA could suppress the pathological changes in spermatozoa caused by HFD or PA treatment. Additionally, the excessive activation of autophagy was also observed in sperm samples from obese, subfertile male patients.

Infertility is defined as the failure to achieve a clinical pregnancy after 12 months or more of regular unprotected sexual intercourse^{1,2}. Approximately 25–30% of the couples experiencing infertility have male-factor infertility³. Obesity is an acknowledged risk factor for male subfertility^{4–7}. The mechanisms underlying obesity-induced male spermatogenesis deficiency remain unclear; deciphering these molecular mechanisms could be of great therapeutic interest for spermatogenesis impairment.

The mechanisms underlying spermatogenesis impairment induced by obesity are complex. Endocrine disorders⁸⁻¹¹, genetic components¹²⁻¹⁵ and physical or chemical factors^{16,17} are all involved in the development of subfertility caused by obesity. However, accumulating data indicate a potential role of autophagy in spermatogenesis^{18,19}.

Autophagy, or programmed cell death type II, is an evolutionarily conserved process that plays a crucial role in maintaining a series of physiological functions through the formation of a double-membrane vesicle termed the autophagosome followed by subsequent fusion with lysosomes and degradation of the cytosolic components by resident hydrolases^{20,21}. Autophagy activates apoptosis, which could lead to oligozoospermia and infertility^{22–24}. Autophagy also regulates inflammation²⁵, which is closely associated with male spermatogenesis impairment^{26–29}. Moreover, a recent study indicated that autophagy plays a key role in the impairment of spermatogenesis after heat treatment³⁰. Germ cell-specific knockout of autophagy-related gene 7 (*Atg7*) causes male infertility in mice¹⁹. Autophagy was also indicated to be involved in the regulation of sperm survival and motility¹⁸. Autophagy was found to be activated under specific circumstances such as nutrient depletion, growth factor depletion and hypoxia^{31,32}. Previous study indicated that mice fed with high-fat diet (HFD) had reduced hepatic autophagy³³, while others reported that nondiabetic HFD mice had upregulated autophagosome formation³⁴. In addition, autophagy is associated with many HFD-related diseases such as nonalcoholic fatty liver disease³⁵, diabetes³⁶ and atherosclerosis³⁷. However, the role of autophagy in obesity-induced spermatogenesis deficiency still remains unknown.

In this study, we used HFD to induce obesity. Both activator and inhibitor of autophagy were employed in this study to explore the role of autophagy in HFD-induced spermatogenesis deficiency. Rapamycin (RAP), which

¹Reproductive Medicine Center, Renmin Hospital of Wuhan University, Wuhan 430060, China. ²Department of Obstetrics and Gynecology, Renmin Hospital of Wuhan University, Wuhan 430060, China. Correspondence and requests for materials should be addressed to J.Y. (email: dryangqing@hotmail.com)



Figure 1. High-fat diet (HFD) induced male mice spermatogenesis impairment. (A) Serum hormone levels of testosterone (T), oestradiol (E2), follicle stimulating hormone (FSH) and luteinizing hormone (LH) (n = 6). ND is defined as normal diet. (B) Hematoxylin and eosin (HE) staining of testis of the indicated groups. Vacuoles in the testis are marked with arrow. Magnification: x200 Scale bar: $50 \,\mu\text{m}$ (n = 6). (C) Statistical results of testis weight/body weight of the indicated groups (n = 6). (D) Statistical analysis of the diameter of seminiferous tubules in four groups (n = 6). (E) Protein level of autophagy marker BECLIN1 in mice testis (n = 6). Full-length gels are presented in Supplementary Figure S4. Data are expressed as mean \pm standard deviation (SD). *p < 0.05, compared with control group.

inhibits the mammalian target of RAP (mTOR) signalling pathway by binding directly to the mTOR complex, and acts as an inducer of autophagy by activating the initiation stage. Chloroquine (CQ), which reverses autophagy by inhibiting lysosomal acidification, results in lysosome accumulation and autophagy blockade. Then, we investigated the role of autophagy in HFD-induced spermatogenesis deficiency by employing CQ to inhibit autophagy and RAP to activate autophagy in mice. The fertility of obese mice was also investigated after CQ and RAP treatment. 3-methyladenine (3-MA), based on its inhibitory effect on class III phosphatidylinositol-3 kinase (PI3K) activity, is another widely used autophagy inhibitor. Palmitic acid (PA) was used in the *in vitro* study to mimic the conditions *in vivo* as described previously^{38,39}. To exclude the off-target effects of CQ and the possibility that the protection of CQ was secondary to the loss of body weight, both CQ and 3-MA were administered via intratesticular injection or were co-cultured with PA *in vivo* and *in vitro*, respectively. In addition, their effects on the viability and motility of mice sperm were explored in this study. Human semen samples were also collected to examine the potential effects of autophagy on the sperm quality of obese subfertile male patients.

Results

Autophagy is over activated in the testis of mice subjected to HFD. As shown in Fig. 1, the mice fed an HFD had impaired spermatogenesis, as manifested by abnormal serum sex hormone levels (Fig. 1A), decreased testis weight/body weight ratio (Fig. 1C) and pathological histological analysis (Fig. 1B), including atrophied seminiferous tubules (Fig. 1D) and an increased number of vacuoles. Meanwhile, after 8 weeks of the HFD, the mice became obese, and the serum cholesterol level was increased (see Supplementary Figure S1). The blood glucose was also detected. After 8 weeks of HFD feeding, the blood glucose was slightly increased. We also



Figure 2. Intraperitoneal injection of chloroquine (CQ) and rapamycin (RAP) on testis autophagy. (A,B) The protein levels of autophagy related markers in mice testis in indicated groups (n = 6). Full-length gels are presented in Supplementary Figure S4. (C) The presence of autophagosomes (arrow) in Leydig cells of mice testis by transmission electron microscopy (TEM) examination of four groups (n = 6). Data are expressed as mean \pm SD. **p* < 0.05, compared with control group. **p* < 0.05, compared with HFD group.

....

use the homeostasis model assessment of insulin resistance (HOMA-IR) to evaluate insulin resistance and found that there was no difference between the control group and HFD group (see Supplementary Figure S1). The autophagy-related protein BECLIN1 was detected to reflect the level of autophagy in the testis of mice subjected to HFD. As indicated, BECLIN1 was found to increase in the HFD group in a time-dependent manner (Fig. 1E and Supplementary Figure S4).

Autophagy is involved in spermatogenesis deficiency induced by HFD. To investigate the role of autophagy in HFD-induced spermatogenesis impairment, CQ and RAP were injected intraperitoneally to inhibit autophagy and induce autophagy, respectively. As illustrated, autophagy was suppressed in mice with CQ injection, as indicated by decreased protein levels of BECLIN1 and increased LC3II/LC3I, p-mTOR and p62 levels. Conversely, the mice treated with RAP showed increased protein levels of BECLIN1 and LC3II/LC3I and decreased p-mTOR and p62 levels, which implied that autophagy was further promoted (Fig. 2A,B and Supplementary Figure S4). As CQ inhibits the degradation of autophagosomes, autophagosomes were increased in both the CQ and RAP treatment groups (Fig. 2C). Subsequently, spermatogenesis function was assessed. As shown in Fig. 3, CQ treatment attenuated HFD-induced spermatogenesis deficiency, as evidenced by increased size of the testis, an improved testis weight/body weight ratio, increased diameter of the seminiferous tubules, and improved serum hormone levels. Conversely, RAP treatment aggravated all these pathological changes. Taken together, these data demonstrate that autophagy is closely involved in HFD-induced spermatogenesis impairment and that inhibiting autophagy with CQ improved spermatogenesis deficiency induced by HFD.

Intraperitoneal injection of CQ on sperm count, viability and motility. Inconsistent with a previous report showing that obese mice had no differences in total sperm count⁴⁰, we found that an HFD led to a decrease in total sperm count. Moreover, HFD decreased the sperm viability and motility to 43% and 40%, respectively.



Figure 3. Suppressing autophagy by CQ ameliorated HFD-induced spermatogenesis deficiency. (A) Serum hormone levels of T, E2, FSH and LH of the four groups (n = 6). (B) Gross morphology of testis in the indicated groups Scale bar: 2 cm. (C) Statistical results of testis weight/body weight ratio of the indicated groups (n = 10). (D) Statistical analysis of the diameter of seminiferous tubules in four groups (n = 10). (E) HE staining of testis of the indicated groups (n = 10). Vacuoles in the testis were marked with arrow. x200 Scale bar: 50 μ m. Data are expressed as mean \pm SD. **p* < 0.05, compared with control group. **p* < 0.05, compared with HFD group.



Figure 4. Inhibiting autophagy by CQ intraperitoneal injection increased sperm viability and motility in mice subjected to HFD. (A–C) Sperm parameters of mice in the indicated groups (n = 8). Data are expressed as mean \pm SD. *p < 0.05, compared with control group. *p < 0.05, compared with HFD group.

Surprisingly, these pathological changes induced by HFD were attenuated after CQ treatment and, indeed, were even slightly potentiated by the addition of the activator of autophagy (Fig. 4).

Intraperitoneal injection of CO on fertility and adverse pregnancy outcomes. The fact that CQ could improve sperm quality in mice with HFD prompted us to explore the effect of CQ on the fertility of obese mice. Inconsistent with a previous study⁴⁰, we found that the percentage of female mice with plugs did not change between the control and HFD groups (Fig. 5A). However, subsequent analysis showed that HFD decreased the pregnancy rate (68.0% vs 32.0%, p = 0.0003). As expected, the pregnancy rate increased to 60.0% (p = 0.005) after CQ treatment and decreased to 20.0% (p = 0.171) after RAP treatment (Fig. 5B). Embryo resorption was also observed in the pregnant mice (Fig. 5C). HFD led to an increase in the embryo resorption rate (2.1% vs 13.0%, p < 0.0001), and conversely, CQ treatment reduced the embryo resorption rate (2.9%, p = 0.003). Intra-uterine foetal death (IUFD) in the pregnant female mice mated with lean and obese male mice was counted, but no significant difference was found in the indicated groups (Fig. 5D).

A single intratesticular injection of CQ and 3-MA improved the motility and viability of sperm *in* **vivo**. To exclude the possibility that the protection of CQ against spermatogenesis impairment was secondary to the loss of body weight, CQ was administered via a single intratesticular injection. We decided the time point of intratesticular injection for the reason that after 4 weeks HFD feeding, the serum testosterone level and the testis weight/body weight began to decrease (Fig. 1). The morphological alteration of testis was also observed at 4 weeks after HFD (Fig. 1). To exclude the nonspecific effect of chloroquine and further confirm the role of inhibiting autophagy in spermatogenesis in HFD mice, mice were subjected to another autophagy inhibitor, 3-MA. As shown in Fig. 6, both CQ and 3-MA improved the motility and viability of sperm relative to that in the HFD group.

CO and **3-MA** attenuated the reduction in the percentages of viable and motile spermatozoa caused by PA. Considering the close relationship between elevated serum free fatty acid level and obesity^{41,42}, and to further decipher the effect of autophagy inhibition, sperm were separated and co-cultured with PA *in vitro* to mimic the conditions *in vivo* as described previously^{38,39}. Moreover, we found that the serum PA level and testis PA level were increased in the mice with HFD (see Supplementary Figure S2). Unpredictably, CQ or 3-MA impaired the viability and motility of sperm (see Supplementary Figure S3). However, both CQ and 3-MA clearly attenuated the reduction in viable and motile spermatozoa caused by PA (Fig. 7A–F).

An increased level of BECLIN1 was observed in obese, subfertile male patients. According to World Health Organization guidelines (REF: WHO 2010), the lower reference limits for semen parameters are: sperm concentration $\geq 15 \times 10^6$ spermatozoa/ml, $\geq 32\%$ progressively motile spermatozoa and $\geq 4\%$ morphologically normal spermatozoa. Decreased semen quality in obese and subfertile patients reflects a condition when at least one of the parameters is under the lower reference limit. BECLINI was found to be increased in obese patients with decreased semen quality, but no significant difference was found between normal individuals and obese and fertile patients, whose semen parameters were above the lower reference limits (see Supplementary Table S1 and Fig. 8A,B), suggesting an association with impaired sperm quality.

Discussion

Autophagy has been implicated in the pathogenesis of different diseases such as cancer^{43,44}, liver disease⁴⁵⁻⁴⁷, kidney disease^{48,49}, neurodegenerative disease^{50,51} and cardiovascular disease^{52,53}. Autophagy was also found to be associated with sperm survival⁵⁴⁻⁵⁶, and a recent study also showed that autophagy and apoptosis work synergistically to induce germ cell death during mouse spermatogenesis³⁰. However, there are no related reports about the role of autophagy in HFD-induced male spermatogenesis impairment. The present study is the first to provide insight into the importance of autophagy during HFD-induced spermatogenesis deficiency and find that autophagy is over activated in HFD-induced spermatogenesis deficiency and CQ attenuates spermatogenesis







Figure 6. The effects of single intratesticular injection of CQ and 3-methyladenine (3-MA) on sperm viability and motility. (A,B) Sperm viability and motility after single intratesticular injection of CQ (n = 6). (C,D) Sperm viability and motility after single intratesticular injection of 3-MA (n = 6). Data are expressed as mean \pm SD. **p* < 0.05, compared with control group. **p* < 0.05, compared with HFD group.

impairment in HFD male mice. CQ also improves sperm count, viability and motility in mice subjected to HFD and decreases adverse pregnancy outcomes. In addition, both *in vivo* and *in vitro* studies demonstrated that CQ and 3-MA improved the reduction in sperm viability and motility induced by HFD and PA treatment, respectively. Moreover, the increased protein level of BECLIN1 in human semen samples, indicated that autophagy was activated in obese, subfertile male patients.

In this study, we showed that mice fed an HFD exhibited impaired spermatogenesis function in a timedependent manner. Meanwhile, the serum steroid hormone levels were obviously changed, as demonstrated by an increased oestradiol (E2) level and decreased testosterone (T), follicle stimulating hormone (FSH) and luteinizing hormone (LH) levels. In addition, these changes were consistent with those of a previous study, which found that seminiferous tubule degeneration is associated with decreased FSH and LH levels⁵⁷. The fertility of HFD male mice was also found to be damaged, which was in accordance with previous studies indicating that a hypercholesterolemic diet contributed to decreased semen quality and reproductive function in male rabbits^{58,59}. Consistent with the previous finding that autophagy was increased in the testis of mice subjected to heat stress and exposed



Figure 7. Suppressing autophagy improved palmitic acid (PA)-induced reduction of sperm viability and motility. (A) mRNA levels of *Beclin1* with CQ treatment (n = 6). (B,C) The effects of CQ on sperm viability and motility *in vivo* (n = 6). (D) mRNA levels of *Beclin1* with 3-MA treatment (n = 6). (E,F) The effects of 3-MA on sperm viability and motility *in vivo* (n = 6). Data are expressed as mean \pm SD. *p < 0.05, compared with control group. *p < 0.05, compared with PA group.

to formaldehyde^{30,60}, we also found that autophagy was over activated in the testis of HFD mice, implying that autophagy may play a role in HFD-induced spermatogenesis deficiency. Previous study indicated that autophagy was activated in human spermatozoa and was involved in the regulation of cell survival and motility¹⁸. Consistent with this finding, the data in our study demonstrated that BECLIN1 was increased in semen samples from obese, subfertile male patients, implying a role of autophagy in human sperm production.

To the best of our knowledge, the primary role of autophagy is as a reparative and cellular cleaning process^{61,62}. The role of autophagy in many other HFD-related diseases has been discussed in the past few decades. Autophagy markers were increased by HFD feeding in many reports^{36,63}, which is consistent with our results, but there is still some evidence supporting a decreased autophagy level with HFD treatment^{36,64}, and dual effects of autophagy in HFD mice were proposed. Recent studies have demonstrated that multiple forms of metabolic stress, including HFD and diabetes, provoke an increase in autophagic flux and that suppression of excessive autophagy could reduce or even reverse the maladaptive response^{65,66}. Moreover, *Atg7* knockdown conferred protection against the heat-induced apoptosis of germ cells³⁰. Consistent with these studies, our study demonstrated that CQ, by inhibiting autophagy, obviously attenuated HFD-induced spermatogenesis deficiency and infertility. Conversely, the mice treated with RAP displayed greater impairments in reproductive function. We also injected CQ and 3-MA directly into the testis and found that an improvement in spermatogenesis was undoubtedly produced by inhibiting autophagy. Taken together, these data demonstrated that inhibition of excessive autophagy could protect against HFD-induced spermatogenesis deficiency and subfertility.

The finding that CQ could improve impaired spermatogenesis does not support the notion that inhibiting autophagy is also a protective mechanism for spermatogenesis at basal condition. Conversely, mitophagy prevents paternal mitochondrial DNA transmission⁶⁷. *Atg7* is required for acrosome biogenesis during spermatogenesis, and germ cell-specific *Atg7*-knockout mice display a phenotype similar to human globozoospermia¹⁹. It was also reported that the activation of autophagy is indispensable for the process of decidualization, which was confirmed in diet-induced obesity female mice⁶⁸. Consistent with these facts, we also found that CQ impaired the viability and motility of sperm *in vitro* and *in vivo* at baseline, implying that autophagy is also indispensable for the physiological process, the results differ. Zhang *et al.* reported that autophagy was not a protective mechanism against heat-induced apoptosis³⁰. Our data also demonstrated that CQ could improve the quality of sperm in mice fed an HFD by inhibiting autophagy. These discrepant results possibly reflect the different roles of autophagy in physiological and pathological processes. Indeed, the dual nature of autophagy is a recurring theme in other organ systems and disease states³¹. Therefore, further studies are required to elucidate the dual roles of autophagy in the production of sperm.





.....

There are several types of cells in testis, including Leydig cells, Sertoli cells and spermatogenic cells. The role of autophagy in these cells during HFD-induced impaired spermatogenesis is still unknown. The finding in our study that CQ or 3-MA could attenuate PA-induced sperm injury imply that CQ or 3-MA may directly target on the sperm. Consistent with a previous study⁶⁹, we also observe the autophagosomes in the Leydig cells, and the decreased T level induced by HFD was restored after CQ treatment, indicating that Leydig cells may mediate CQ-induced protection. The results that CQ or RAP can not affect the level of LH and FSH in the mice with HFD, implying that pituitary gland was not involved in the alteration of hormones caused by the treatment of CQ or RAP.

Despite the results that autophagy was over activated in spermatogenesis when males were exposed to an HFD, we do not yet know which molecules induced such activation. Indeed, an HFD induced the decreased phosphorylation of adenosine monophosphate-activated protein kinase (AMPK α)³⁹ and thus inhibited the activation of mTOR, which is an agonist of autophagy^{70,71}. Previous reports also demonstrated that autophagy is activated in response to different forms of metabolic stress, including nutrient deprivation, growth factor depletion, and hypoxia^{31,32}. Whether these metabolic stresses play roles in HFD-induced spermatogenesis impairment remains unknown. Genetic approaches to inhibit autophagy will yield more conclusive information about the biologic roles of autophagy in spermatogenesis and HFD-induced spermatogenesis deficiency. In addition, our study does not decipher whether HFD-induced autophagy occurs in response to the inciting stress, is a secondary response, promotes or antagonizes disease, or exists as an epiphenomenon.

In conclusion, we have confirmed that autophagy is over activated in HFD-induced spermatogenesis deficiency and found that inhibiting the autophagic process ameliorates HFD-induced spermatogenesis impairment *in vitro* and *in vivo*. Our study also indicate that autophagy is increased in semen samples from obese, subfertile male patients. Taken together, our finding provides a new therapeutic target for infertility induced by obesity, and further investigations are required to clarify the specific molecular mechanism and signalling pathway that are involved in the activation of autophagy, especially in human subjects.

Methods

Reagents. CQ (100421-200401) was purchased from the National Institutes for Food and Drug Control (Beijing, China), and RAP (R-5000) was purchased from LC Laboratories (Woburn, MA, USA). 3-MA (M9281) was purchased from Sigma (St. Louis, MO, USA). The rabbit anti-SQSTM1/p62 monoclonal antibody (ab109012) for western blot was purchased from Abcam (Cambridge, UK). The rabbit anti-phospho-mTOR antibody (#2974), rabbit anti-mTOR antibody (#2983), rabbit anti-BECLIN1 antibody (#3495) and rabbit anti-LC3II/I antibody (#12741) for western blot were purchased from Cell Signaling Technology (Beverly, MA, USA). The rabbit anti-GAPDH antibody (sc-25778) for western blot was purchased from Santa Cruz Biotechnology (Dallas, TX, USA). The secondary antibody was purchased from LI-COR Biosciences. Oestradiol (E2, E-EL-0065c), testosterone (T, E-EL-0072c), follicle stimulating hormone (FSH, E-EL-M0511c), and luteinizing hormone (LH, E-EL-M0057c) detection kits were purchased from Elabscience Biotechnology Co., Ltd. (Wuhan, China). PA (P9767) was purchased from Sigma (St. Louis, MO, USA). All the other chemicals used in our study were of analytical grade.

Animal care and treatment. All the animal experiments in our study were performed in accordance with the Guidelines for the Care and Use of Laboratory Animals published by the United States National Institutes of Health (NIH Publication, revised 2011) and the Guidelines for the Care and Use of Laboratory Animals of the Chinese Animal Welfare Committee and were approved by the Animal Use Committees of Renmin Hospital of Wuhan University. Male C57BL mice (Permit number: 42000600004159, body weight: 20-22 g) were obtained from Hubei Provincial Center for Disease Control and Prevention (Wuhan, China). The mice were housed under a 12h light: 12h dark cycle in a temperature (20-25 °C) and humidity ($50 \pm 5\%$)-controlled environment with free access to food and water. Mice (n = 40) were randomly divided into four groups, including a control group (Normal diet + 0.1% DMSO, n = 10), an HFD group (HFD + 0.1% DMSO, n = 10), an HFD + CQ group (n = 10) and an HFD + RAP (n = 10) group. All the mice were fed ad libitum either a standard chow or a HFD. Subsequently, the mice in the HFD + CQ or HFD + RAP groups were intraperitoneally injected with CQ $(10 \text{ mg/ kg})^{72}$ or RAP $(1 \text{ mg/ kg})^{73}$ once per day for 8 weeks^{74,75}, and the mice in the control group were given the same volume of vehicle. CQ was dissolved in sterile saline, and RAP was dissolved in 0.1% DMSO. After that, the mice were anaesthetized with an intraperitoneal injection of 3% sodium pentobarbital, and blood samples were obtained from the angular veins. Then, mice were euthanized with an intraperitoneal injection excessive sodium pentobarbital, the testes were collected for further experiments and the testis weight/body weight ratio was calculated. To further confirm the effect of autophagy inhibition on spermatogenesis in HFD male mice, a single intratesticular injection was performed. After mice had received the HFD for 4 weeks, $40 \,\mu$ l of the autophagy inhibitor CQ ($300 \,\mu$ M) or 3-MA $(10 \text{ mM}, \text{dissolved in sterile saline})^{19}$ was injected under the tunica vaginalis of the testes with a 50 µl micro-injector at a depth of 5 mm under the tunica albuginea as described previously⁷⁶. After injection, the mice were fostered for another 4 weeks. Spermatozoa were taken from the epididymis to examine their viability and motility.

Determination of serum hormone levels. Blood samples were collected from angular veins. After centrifuging the samples at 1409 g and 4 °C for 15 minutes, the supernatant sera were obtained and stored at -80 °C until analysis. The levels of T, E2, FSH and LH in the serum were measured using commercially available ELISA kits according to the manufacturer's instructions. Briefly, ninety-six-well plates were coated with the primary antibody at 37 °C for 2h. After being blocked with 5% milk, the serum were added into the plates. The plates were incubated at 37 °C for 2h. After that, the plates were incubated with peroxidase-conjugated second antibodies at 37 °C for 1h. After that, the optical density values were detected by a microplate reader (Biotek, Vermont, USA).

Histological analysis. The sections of testis tissue were routinely fixed in Bouin's solution, dehydrated, and embedded in paraffin. Testis tissue sections were stained with haematoxylin and eosin (H&E) to examine morphology. Sections (5μ m) were viewed under light microscopy (Nikon E100), and photomicrographs were captured with the Photo Imaging System (Canon 600D). The diameter of seminiferous tubules was determined using Image-Pro Plus 6.0. In each group, 100 seminiferous tubules (5 fields per mouse, 2 random seminiferous tubules per field) were measured in 10 mice, and the mean seminiferous tubule diameter was determined.

Protein extraction and western blot assay. RIPA buffer (720 µl radioimmunoprecipitation buffer, 100 mmol/l PMSF, 100 µl cocktail, 100 µl Phos-stop, 20 mmol/l NaF and 100 mmol/l Na₃VO₄ in 1 ml) was used to extract the proteins from testis tissue of mice and sperm samples of mice and humans⁷⁷. The protein concentrations were detected with a BCA Protein Assay Kit (Pierce, Rockford, IL, USA). Proteins (20 µg) were separated on SDS-PAGE gels (LC31/II: 15%; mTOR and p-mTOR: 6%; Others: 10%) for 2 h (75 V 30 min; 100 V 90 min) at room temperature and then transferred to a PVDF membrane (IPVH00010, Millipore, Billerica, MA, USA) at 4°C for different times (200 mA; LC31/II: 70 min; mTOR and p-mTOR: 120 min; Others: 90 min). After being blocked with 5% non-fat milk, the membranes were incubated overnight at 4°C with the following primary antibodies: rabbit anti-GAPDH antibody (1:5000), rabbit anti-BECLIN1 antibody (1:1000), rabbit anti-SQSTM/p62 monoclonal antibody (1:1000), rabbit anti-LC3II/I antibody (1:1000), rabbit anti-Phospho-mTOR antibody (1:1000) and rabbit anti-mTOR antibody (1:1000). After incubation with the secondary antibody (1:5000), the membranes were scanned using a two-colour infrared imaging system (Odyssey, LI-COR, Lincoln, NE, USA). Densitometry analysis was performed by Odyssey as described previously⁷⁸, and the results were normalized to GAPDH.

RNA isolation and qPCR analysis of mouse semen samples. Sperm samples of mice were homogenized in TRIzol (Invitrogen, Massachusetts, USA) on ice, and total mRNA was extracted. mRNA was reverse transcribed into cDNA using the Transcriptor First Strand cDNA Synthesis Kit (04896866001, Roche, USA) according to the manufacturer's instructions. All experiments were performed in triplicate, and quantitative RT-PCR analysis was carried out using the Light Cycler 480 SYBR Green 1 Master Mix (04707516001, Roche, USA). The primer sequences for *Beclin1* were: forward: 5'-ATCCTGGACCGTGTCACCATCCAGG-3'; reverse: 5'-GTTGAGCTGAGTGTCCAGCTGG-3'. The primer sequences for *Gapdh* were: forward: 5'-GTTGTCTCCTGCGACTTCA-3'; reverse: 5'-GGTGGTCCAGGGTTTCTTA-3'. The mRNA levels were analysed using the $2^{-\Delta \Delta Ct}$ method and normalized to *Gapdh*. The qPCR analysis was performed as described in a previous report⁷⁹.

Transmission electron microscopy (TEM). Testis tissue was fixed with 2.5% glutaraldehyde, and then these tissues were post-fixed in 1% OsO_4 for 2 h. After rehydration, these tissues were embedded in epoxy resin. Ultrathin sections (80 nm) were collected on formvar-coated copper grids. Subsequently, these grids were counter-stained with uranyl acetate and lead citrate and examined on an EM2010FEF- Ω transmission electron microscope (JEOL, Tokyo, Japan). Autophagosomes were detected as described in a previous study⁸⁰ and identified by the presence of a double-membrane vesicle that contained cytosol and/or organelles and looked morphologically intact.

Semen analysis in mice. The epididymis was carefully dissected away from the fat. The separated epididymides were immediately placed into Ringer's solution and dissected to determine the count, viability and motility of the sperm as described in a previous study⁴⁰. The sperm were gradually pressed out from the epididymis and collected in an Eppendorf tube. The sperm number was counted with a haemocytometer independently for three times.

Eosin-nigrosin staining solution (1% eosin and 10% nigrosin) was used to detect sperm viability. Semen samples were first incubated with the eosin-nigrosin staining solution and smeared on a microscope slide. The spermatozoa were observed under a light microscope at 100x magnification. Live spermatozoa is not stained, while dead spermatozoa with disintegrating cell membranes take up the stain⁸¹. Sperm motility was tested with computer-assisted sperm analysis (CASA). Sperm were incubated in Ringer's solution at room temperature for 30 minutes and placed into CASA assay chambers (Hamilton Thorne Research, Beverly, MA, USA). Sperm tracks (1.5 s, 30 frames) were captured at 60 Hz and were analysed with HTM-IVOS Sperm Analyzer software (version 12.2 L; Hamilton Thorne)⁸². Five hundred sperm cells were randomly selected and evaluated to obtain the sperm viability and sperm motility; at least three replications were completed for each sample.

Assessment of the fertility of male mice. At the end of the 8-week HFD, the mice receiving different treatments were used in a breeding assay. Each male mouse (n = 5) was caged with two females for five days; consequently, a total of 50 females were involved in the study, and their vaginal plugs were checked every morning. The day on which the vaginal plug observed was defined as day 0.5 of gestation, and on day 14.5 of gestation, female mice were euthanized to count the number of dead foetuses and resorptions.

In vitro study. Mouse sperm samples obtained from the cauda epididymis were diluted to attain a concentration of 5×10^6 spermatozoa/ml and incubated in G-IVF (Vitrolife, Sweden) medium, fatty medium (1 mM PA)⁸³ or fatty medium with CQ (100 μ M) and 3-MA (10 mM)⁸⁴. Incubations were performed under a humidified atmosphere of 5% CO₂/95% O₂ and 37 °C for 5 h, and sperm viability and motility were assessed as described above. Three independent experiments were carried out in the *in vitro* study.

Human studies. The human studies conformed to the Declaration of Helsinki and were approved by the Human Research Ethics Committees of Renmin Hospital of Wuhan University in Wuhan, China. All the participants in this study were interviewed after signing an informed consent. Semen samples were obtained from subjects who attended the infertility clinic by masturbation after 3-7 days of sexual abstinence. The interviewers collected the data of all participants, including their ethnicity, age, height, body weight, medical history, and lifestyle through a structured questionnaire. The subjects were ethnic Han Chinese from Hubei province and its nearby regions. They were men who visited the Reproductive Medical Center of Renmin Hospital of Wuhan University for infertility treatment from June 2014 to December 2014. All subjects were age-matched, and men who were unhealthy with other causes of defective spermatogenesi⁸⁵, including infection, varicocele, obstruction of the vas deferens, chromosomal abnormalities or smokers, were excluded. Finally, all semen samples were divided into three groups, including the control group (n = 85), obese and fertile group (n = 65), and obese and subfertile group (n = 79). The semen samples in the control group were from individuals with normal body mass index (BMI) and normal semen parameters as described above. The semen samples in the obese and fertile group were from men whose BMIs were more than 30 and had normal semen quality, while the semen samples in the obese and subfertile group were from those whose BMIs were more than 30 and had decreased semen quality. The semen samples were liquefied for 15–30 minutes at 37 °C, and subsequently the semen parameters were tested in accordance with the World Health Organization guidelines. The residual semen samples were used for protein extraction and subsequent western blotting analysis.

Data analysis. Group data are reported as the mean \pm standard deviation (SD). All statistical tests were performed with SPSS 19.0. All the data in the present study were normally distributed (p < 0.05), as determined by the Kolmogorov-Smirnov's test. Two-group comparisons were analysed with an unpaired Student's t-test after confirming equality of variance with the F test. Multiple-group comparisons were performed using one-way ANOVA followed by Tukey's post hoc test⁷⁰ when ANOVA analysis indicated a significant value of F and no variance in homogeneity; otherwise, Tamhane's T2 post hoc test was used. The percentage of plugs, the pregnancy rate, the resorption rate and the IUFD between different groups were compared using a chi-square test. *P*-values less than 0.05 were considered significant.

References

- 1. Chandra, A., Martinez, G. M., Mosher, W. D., Abma, J. C. & Jones, J. Fertility, family planning, and reproductive health of US women: data from the 2002 National Survey of Family Growth. *Vital Health Stat* 23, 1–160 (2005).
- 2. Chandra, A. & Stephen, E. H. Infertility service use among US women: 1995 and 2002. FERTIL STERIL 93, 725-736 (2010).
- Hammoud, A. O., Gibson, M., Peterson, C. M., Hamilton, B. D. & Carrell, D. T. Obesity and male reproductive potential. J Androl 27, 619–626 (2006).
- Jensen, T. K. et al. Body mass index in relation to semen quality and reproductive hormones among 1,558 Danish men. FERTIL STERIL 82, 863–870 (2004).
- 5. Kort, H. I. et al. Impact of body mass index values on sperm quantity and quality. J Androl 27, 450-452 (2006).
- Bieniek, J. M. et al. Influence of increasing body mass index on semen and reproductive hormonal parameters in a multi-institutional cohort of subfertile men. FERTIL STERIL 106, 1070–1075 (2016).
- Andersen, J. M. et al. Body Mass Index Is Associated with Impaired Semen Characteristics and Reduced Levels of Anti-Mullerian Hormone across a Wide Weight Range. PLOS ONE 10, e130210 (2015).
- Fui, M. N., Dupuis, P. & Grossmann, M. Lowered testosterone in male obesity: mechanisms, morbidity and management. ASIAN J ANDROL 16, 223–231 (2014).
- Camacho, E. M. *et al.* Age-associated changes in hypothalamic-pituitary-testicular function in middle-aged and older men are modified by weight change and lifestyle factors: longitudinal results from the European Male Ageing Study. *EUR J ENDOCRINOL* 168, 445–455 (2013).
- Tajar, A. et al. Characteristics of secondary, primary, and compensated hypogonadism in aging men: evidence from the European Male Ageing Study. J Clin Endocrinol Metab 95, 1810–1818 (2010).
- 11. Ayanian, S. & Irwig, M. S. Hypogonadism in a male-to-female transsexual with super obesity. ANDROLOGIA 45, 285-288 (2013).
- Aksglaede, L. et al. Serum concentrations of Anti-Mullerian Hormone (AMH) in 95 patients with Klinefelter syndrome with or without cryptorchidism. ACTA PAEDIATR 100, 839–845 (2011).
- 13. Dupont, C. et al. Obesity leads to higher risk of sperm DNA damage in infertile patients. ASIAN J ANDROL 15, 622-625 (2013).
- Hammoud, A. *et al.* An aromatase polymorphism modulates the relationship between weight and estradiol levels in obese men. *FERTIL STERIL* 94, 1734–1738 (2010).
- Chavarro, J. E., Toth, T. L., Wright, D. L., Meeker, J. D. & Hauser, R. Body mass index in relation to semen quality, sperm DNA integrity, and serum reproductive hormone levels among men attending an infertility clinic. *FERTIL STERIL* 93, 2222–2231 (2010).
 Hammoud, A. O. *et al.* Obesity and male infertility: a practical approach. *SEMIN REPROD MED* 30, 486–495 (2012).
- Hammoud, A. O. *et al.* Obesity and materinity: a practical approach. *JEMIN REPROD MED* 30, 486–495 (2012).
 Hammoud, A. O., Carrell, D. T., Gibson, M., Peterson, C. M. & Meikle, A. W. Updates on the relation of weight excess and
- reproductive function in men: sleep apnea as a new area of interest. ASIAN J ANDROL 14, 77-81 (2012).
- 18. Aparicio, I. M. *et al.* Autophagy-related proteins are functionally active in human spermatozoa and may be involved in the regulation of cell survival and motility. *Sci Rep* **6**, 33647 (2016).
- 19. Wang, H. et al. Atg7 is required for acrosome biogenesis during spermatogenesis in mice. CELL RES 24, 852-869 (2014).
- 20. Yang, Z. & Klionsky, D. J. Eaten alive: a history of macroautophagy. NAT CELL BIOL 12, 814–822 (2010).
- 21. Weide, T. & Huber, T. B. Implications of autophagy for glomerular aging and disease. CELL TISSUE RES 343, 467-473 (2011).
- Rotter, V. et al. Mice with reduced levels of p53 protein exhibit the testicular giant-cell degenerative syndrome. Proc Natl Acad Sci USA 90, 9075–9079 (1993).
- Knudson, C. M., Tung, K. S., Tourtellotte, W. G., Brown, G. A. & Korsmeyer, S. J. Bax-deficient mice with lymphoid hyperplasia and male germ cell death. SCIENCE 270, 96–99 (1995).
- 24. Lee, H. W. et al. Essential role of mouse telomerase in highly proliferative organs. NATURE **392**, 569–574 (1998).
- 25. Levine, B., Mizushima, N. & Virgin, H. W. Autophagy in immunity and inflammation. NATURE 469, 323-335 (2011).
- Baumber, J., Sabeur, K., Vo, A. & Ball, B. A. Reactive oxygen species promote tyrosine phosphorylation and capacitation in equine spermatozoa. *THERIOGENOLOGY* 60, 1239–1247 (2003).
- Baccetti, B., Collodel, G. & Piomboni, P. Apoptosis in human ejaculated sperm cells (notulae seminologicae 9). J Submicrosc Cytol Pathol 28, 587–596 (1996).
- 28. Oosterhuis, G. J. *et al.* Measuring apoptosis in human spermatozoa: a biological assay for semen quality? *FERTIL STERIL* 74, 245–250 (2000).
- Haidl, G., Allam, J. P. & Schuppe, H. C. Chronic epididymitis: impact on semen parameters and therapeutic options. ANDROLOGIA 40, 92–96 (2008).
- 30. Zhang, M. *et al.* Autophagy and apoptosis act as partners to induce germ cell death after heat stress in mice. *PLOS ONE* 7, e41412 (2012).
- 31. Araya, J., Hara, H. & Kuwano, K. Autophagy in the pathogenesis of pulmonary disease. Intern Med 52, 2295–2303 (2013).
- Malicdan, M. C., Noguchi, S. & Nishino, I. Autophagy in a mouse model of distal myopathy with rimmed vacuoles or hereditary inclusion body myopathy. AUTOPHAGY 3, 396–398 (2007).
- Liu, H. Y. et al. Hepatic autophagy is suppressed in the presence of insulin resistance and hyperinsulinemia: inhibition of FoxO1dependent expression of key autophagy genes by insulin. J BIOL CHEM 284, 31484–31492 (2009).
- 34. Ebato, C. et al. Autophagy is important in islet homeostasis and compensatory increase of beta cell mass in response to high-fat diet. CELL METAB 8, 325–332 (2008).
- Mao, Y. et al. Ghrelin Attenuated Lipotoxicity via Autophagy Induction and Nuclear Factor-kappaB Inhibition. CELL PHYSIOL BIOCHEM 37, 563–576 (2015).
- 36. Liu, Y. *et al.* Adiponectin stimulates autophagy and reduces oxidative stress to enhance insulin sensitivity during high-fat diet feeding in mice. *DIABETES* 64, 36–48 (2015).
- Wang, B., Zhong, Y., Huang, D. & Li, J. Macrophage autophagy regulated by miR-384-5p-mediated control of Beclin-1 plays a role in the development of atherosclerosis. AM J TRANSL RES 8, 606–614 (2016).
- 38. Zeng, W. et al. Inhibition of HMGB1 release via salvianolic acid B-mediated SIRT1 up-regulation protects rats against non-alcoholic fatty liver disease. Sci Rep 5, 16013 (2015).
- Kuwabara, Y. et al. MicroRNA-451 exacerbates lipotoxicity in cardiac myocytes and high-fat diet-induced cardiac hypertrophy in mice through suppression of the LKB1/AMPK pathway. CIRC RES 116, 279–288 (2015).
- Ghanayem, B. I., Bai, R., Kissling, G. E., Travlos, G. & Hoffler, U. Diet-induced obesity in male mice is associated with reduced fertility and potentiation of acrylamide-induced reproductive toxicity. BIOL REPROD 82, 96–104 (2010).
- Groop, L. C. et al. Glucose and free fatty acid metabolism in non-insulin-dependent diabetes mellitus. Evidence for multiple sites of insulin resistance. J CLIN INVEST 84, 205–213 (1989).
- 42. Manco, M. *et al.* Insulin resistance directly correlates with increased saturated fatty acids in skeletal muscle triglycerides. *METABOLISM* **49**, 220–224 (2000).
- 43. Guo, J. Y., Xia, B. & White, E. Autophagy-mediated tumor promotion. CELL 155, 1216-1219 (2013).
- 44. Levine, B. Cell biology: autophagy and cancer. NATURE 446, 745-747 (2007).
- Alavian, S. M. *et al.* Virus-triggered autophagy in viral hepatitis possible novel strategies for drug development. J Viral Hepat 18, 821–830 (2011).
- Wang, K., Damjanov, I. & Wan, Y. J. The protective role of pregnane X receptor in lipopolysaccharide/D-galactosamine-induced acute liver injury. LAB INVEST 90, 257–265 (2010).

- Osna, N. A., Thomes, P. G. & Donohue, T. M. Jr. Involvement of autophagy in alcoholic liver injury and hepatitis C pathogenesis. World J Gastroenterol 17, 2507–2514 (2011).
- Asanuma, K. *et al.* MAP-LC3, a promising autophagosomal marker, is processed during the differentiation and recovery of podocytes from PAN nephrosis. *FASEB J* 17, 1165–1167 (2003).
- Chien, C. T., Shyue, S. K. & Lai, M. K. Bcl-xL augmentation potentially reduces ischemia/reperfusion induced proximal and distal tubular apoptosis and autophagy. TRANSPLANTATION 84, 1183–1190 (2007).
- Banerjee, R., Beal, M. F. & Thomas, B. Autophagy in neurodegenerative disorders: pathogenic roles and therapeutic implications. TRENDS NEUROSCI 33, 541–549 (2010).
- 51. Hara, T. et al. Suppression of basal autophagy in neural cells causes neurodegenerative disease in mice. NATURE 441, 885–889 (2006).
- 52. Marzetti, E. *et al.* Role of mitochondrial dysfunction and altered autophagy in cardiovascular aging and disease: from mechanisms to therapeutics. *Am J Physiol Heart Circ Physiol* **305**, H459–H476 (2013).
- Nemchenko, A., Chiong, M., Turer, A., Lavandero, S. & Hill, J. A. Autophagy as a therapeutic target in cardiovascular disease. J MOL CELL CARDIOL 51, 584–593 (2011).
- 54. Gallardo, B. J. et al. Autophagy and apoptosis have a role in the survival or death of stallion spermatozoa during conservation in refrigeration. PLOS ONE 7, e30688 (2012).
- Bolanos, J. M. et al. During cooled storage the extender influences processed autophagy marker light chain 3 (LC3B) of stallion spermatozoa. ANIM REPROD SCI 145, 40–46 (2014).
- Aparicio, I. M., Martin, M. P., Salido, G. M., Pena, F. J. & Tapia, J. A. The autophagy-related protein LC3 is processed in stallion spermatozoa during short-and long-term storage and the related stressful conditions. ANIMAL 10, 1182–1191 (2016).
- Sharma, S. *et al.* Free fatty acids induce Lhb mRNA but suppress Fshb mRNA in pituitary LbetaT2 gonadotropes and diet-induced obesity reduces FSH levels in male mice and disrupts the proestrous LH/FSH surge in female mice. *ENDOCRINOLOGY* 154, 2188–2199 (2013).
- 58. Saez, L. T. *et al.* Semen quality and sperm function loss by hypercholesterolemic diet was recovered by addition of olive oil to diet in rabbit. *PLOS ONE* **8**, e52386 (2013).
- 59. Saez, L. T. et al. Hypercholesterolemia impaired sperm functionality in rabbits. PLOS ONE 5, e13457 (2010).
- 60. Han, S. P. *et al.* Formaldehyde exposure induces autophagy in testicular tissues of adult male rats. *ENVIRON TOXICOL* **30**, 323–331 (2015).
- Levine, B. & Klionsky, D. J. Development by self-digestion: molecular mechanisms and biological functions of autophagy. DEV CELL 6, 463–477 (2004).
- Rodrigues, P., Limback, D., McGinnis, L. K., Plancha, C. E. & Albertini, D. F. Multiple mechanisms of germ cell loss in the perinatal mouse ovary. *REPRODUCTION* 137, 709–720 (2009).
- 63. Sun, Q. et al. Factors that Affect Pancreatic Islet Cell Autophagy in Adult Rats: Evaluation of a Calorie-Restricted Diet and a High-Fat Diet. PLOS ONE 11, e151104 (2016).
- 64. Codogno, P. & Meijer, A. J. Autophagy: a potential link between obesity and insulin resistance. CELL METAB 11, 449-451 (2010).
- 65. Zhang, T. Z. et al. Suppressing autophagy protects photoreceptor cells from light-induced injury. Biochem Biophys Res Commun 450, 966–972 (2014).
- 66. Ojha, R., Jha, V., Singh, S. K. & Bhattacharyya, S. Autophagy inhibition suppresses the tumorigenic potential of cancer stem cell enriched side population in bladder cancer. *Biochim Biophys Acta* 1842, 2073–2086 (2014).
- 67. Al, R. S. *et al.* Postfertilization autophagy of sperm organelles prevents paternal mitochondrial DNA transmission. *SCIENCE* **334**, 1144–1147 (2011).
- 68. Rhee, J. S. *et al.* Diet-induced obesity impairs endometrial stromal cell decidualization: a potential role for impaired autophagy. *HUM REPROD* **31**, 1315–1326 (2016).
- 69. Tang, X. M., Clermont, Y. & Hermo, L. Origin and fate of autophagosomes in Leydig cells of normal adult rats. J Androl 9, 284–293 (1988).
- Ma, Z. G. et al. Asiatic Acid Protects against Cardiac Hypertrophy through Activating AMPKalpha Signalling Pathway. INT J BIOL SCI 12, 861–871 (2016).
- 71. Ma, Z. G. *et al.* Protection against cardiac hypertrophy by geniposide involves the GLP-1 receptor/AMPKalpha signalling pathway. *Br J Pharmacol* **173**, 1502–1516 (2016).
- Okanlawon, A. O., Noronha, C. C. & Ashiru, O. A. An investigation into the effects of chloroquine on fertility of male rats. West Afr J Med 12, 118–121 (1993).
- Scarpace, P. J. et al. Rapamycin Normalizes Serum Leptin by Alleviating Obesity and Reducing Leptin Synthesis in Aged Rats. J Gerontol A Biol Sci Med Sci 71, 891–899 (2016).
- 74. Liu, Z. *et al.* Vitamin B6 Prevents Endothelial Dysfunction, Insulin Resistance, and Hepatic Lipid Accumulation in Apoe (-/-) Mice Fed with High-Fat Diet. J DIABETES RES 2016, 1748065 (2016).
- 75. Yoo, D. Y. *et al.* Reduced cell proliferation and neuroblast differentiation in the dentate gyrus of high fat diet-fed mice are ameliorated by metformin and glimepiride treatment. *NEUROCHEM RES* **36**, 2401–2408 (2011).
- Sato, M., Ishikawa, A. & Kimura, M. Direct injection of foreign DNA into mouse testis as a possible *in vivo* gene transfer system via epididymal spermatozoa. *MOL REPROD DEV* 61, 49–56 (2002).
- 77. Mu, Y., Yan, W. J., Yin, T. L. & Yang, J. Curcumin ameliorates highfat dietinduced spermatogenesis dysfunction. *MOL MED REP* 14, 3588–3594 (2016).
- Liu, Y. et al. Toll-interacting protein (Tollip) negatively regulates pressure overload-induced ventricular hypertrophy in mice. CARDIOVASC RES 101, 87–96 (2014).
- 79. Bustin, S. A. *et al.* The MIQE guidelines: minimum information for publication of quantitative real-time PCR experiments. *CLIN CHEM* **55**, 611–622 (2009).
- 80. Klionsky, D. J. et al. Guidelines for the use and interpretation of assays for monitoring autophagy. AUTOPHAGY 8, 445-544 (2012).
- Kokilavani, P. et al. Antioxidant mediated ameliorative steroidogenesis by Commelina benghalensis L. and Cissus quadrangularis L. against quinalphos induced male reproductive toxicity. Pestic Biochem Physiol 109, 18–33 (2014).
- Wei, Z. T. et al. The long-term effects of superovulation on fertility and sexual behavior of male offspring in mice. J Assist Reprod Genet 31, 555–560 (2014).
- Yin, J. et al. Rapamycin improves palmitate-induced ER stress/NF kappa B pathways associated with stimulating autophagy in adipocytes. Mediators Inflamm 2015, 272313 (2015).
- Teng, A. C. *et al.* Metformin increases degradation of phospholamban via autophagy in cardiomyocytes. *Proc Natl Acad Sci USA* 112, 7165–7170 (2015).
- 85. Yu, B. *et al.* Cigarette smoking is associated with abnormal histone-to-protamine transition in human sperm. *FERTIL STERIL* **101**, 51–57 (2014).

Acknowledgements

This work was supported by the National Natural Science Foundation of China (81370767, grant recipient J.Y.), the National Natural Science Foundation of China (81401255, grant recipient J.L.) and the Natural Science Foundation of Hubei Province (2015CFB722, grant recipient Y.Z.).

Author Contributions

Y.M., W.J.Y., T.L.Y., Y.Z. and J.Y. were involved in the study design and preparation of the manuscript. Y.M. carried out the experiments in the study with W.J.Y., T.L.Y. and J.L. analyzed the data, drafted the manuscript and had critical discussion with Y.Z., J.Y. and W.J.Y.

Additional Information

Supplementary information accompanies this paper at http://www.nature.com/srep

Competing Interests: The authors declare no competing financial interests.

How to cite this article: Mu, Y. *et al.* Diet-induced obesity impairs spermatogenesis: a potential role for autophagy. *Sci. Rep.* 7, 43475; doi: 10.1038/srep43475 (2017).

Publisher's note: Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit http://creativecommons.org/licenses/by/4.0/

© The Author(s) 2017